

A.Kh. Attaev\*

*Institute of Applied Mathematics and Automation of Kabardin-Balkar Scientific Center of RAS, Nalchik, Russia  
(E-mail: [attaev.anatoly@yandex.ru](mailto:attaev.anatoly@yandex.ru))*

## On Some Non-local Boundary Value and Internal Boundary Value Problems for the String Oscillation Equation

The work is devoted to the problem of setting new boundary and internal boundary value problems for hyperbolic equations. The consideration of these settings is given on the example of a wave equation. The research involves the d'Alembert method, the mean value theorem and the method of successive approximations. The paper formulates and studies a number of non-local problems summarizing the classical Goursat and Darbu tasks. Some of them are marginal, and the other part is internal-marginal, and in both cases both characteristic and uncharacteristic displacements are considered. It should also be noted that a number of problems discussed below arose as a special case in the construction of the theory of correct problems for the model loaded equation of string oscillation.

*Keywords:* Wave equation, general solution, Cauchy problem, Goursat problem, Darboux problem, problem with characteristic shift, problem with uncharacteristic displacement.

### *Introduction*

Boundary value problems for partial differential equations with nonlocal conditions present a class of problems solvability of which is important both for differential equations development and for their applications. They can be used in mathematical modeling as well as in the theory of loaded equations and coefficient inverse problems.

In 1969, in the work [1] A.M. Nakhushhev proposed a number of problems of a new type, which entered the mathematical literature under the name of problems with displacement. These tasks were announced as part of the implementation of the problem of finding correctly posed problems for second-order mixed-type equations with two independent variables, put forward in the 60s of the last century by A.V. Bitsadze.

In accordance with the classification proposed by him in [2, 3], these problems, bounded by two intersecting characteristics of a given hyperbolic equation and one characteristic line, are non-local and with an edge offset. The bibliography of works devoted to regular local and nonlocal boundary value problems for strictly hyperbolic equations is very extensive and is most fully given in the monograph [4]. Let us note some of them that are closest to the problems discussed in this paper [1–11].

In this paper, a number of non-local problems generalizing the classical problems of Goursat and Darboux are formulated and investigated. Some of them are marginal, and the other part is internally marginal, and in both cases both characteristic and non-characteristic displacements are considered. It should also be noted that a number of the problems discussed below arose as a special case when constructing the theory of correct problems by analogy with [7] for the model loaded string oscillation equation.

---

\*Corresponding author.

*E-mail: [attaev.anatoly@yandex.ru](mailto:attaev.anatoly@yandex.ru)*

1 The main part

In this paper, the wave equation is considered as a model equation with second-order partial derivatives of two independent variables  $x$  and  $y$  a hyperbolic equation

$$u_{xx} - u_{yy} = 0. \tag{1}$$

Let  $\Omega_1$  be a simply connected domain of the plane of a complex variable  $z = x + iy$ , bounded by the characteristics  $AB, BC, CD$  and  $DA$  equation (1), coming out of the points, respectively  $A(0, 0), B(\frac{1}{2}, \frac{1}{2}), C(1, 0)$  and  $D(\frac{1}{2}, -\frac{1}{2})$ ;  $AC = \bar{J}$ . By  $\Omega_2$  we denote the area bounded by the characteristics  $AB, BC$  and the segment  $AC$  of the straight line  $y = 0$ .

By the regular solution of the equation (1) we will understand any function  $u(x, y) \in C(\bar{\Omega}_i) \cap C^2(\Omega_i)$ ,  $i = 1, 2$  satisfying equation (1).

2 Non-local problems with edge displacement

*Problem 2.1.* Find regular in the area  $\Omega_1$  decision  $u(x, y)$  equation (1), satisfying conditions

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{1+x}{2}, \frac{x-1}{2}\right) = \gamma_1(x), \quad x \in \bar{J},$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{1+x}{2}, \frac{1-x}{2}\right) = \gamma_2(x), \quad x \in \bar{J},$$

where

$$\alpha_1(0) \cdot \alpha_1(1) \neq \alpha_2(0) \cdot \alpha_2(1), \tag{2}$$

$$\alpha_1(x) \cdot \alpha_2(x) \neq 1, \forall x \in \bar{J}, \tag{3}$$

$$\alpha_i(x), \gamma_i(x) \in C(\bar{J}) \cap C^2(J), \quad i = 1, 2.$$

Based on the Asgerisson principle for characteristic quadrilaterals with vertices respectively at point

$$(0, 0), \left(\frac{x+1}{2}, \frac{x-1}{2}\right), \left(\frac{x}{2}, \frac{x}{2}\right), \left(\frac{1}{2}, -\frac{1}{2}\right)$$

and

$$(0, 0), \left(\frac{1+x}{2}, \frac{1-x}{2}\right), \left(\frac{x}{2}, -\frac{x}{2}\right), \left(\frac{1}{2}, \frac{1}{2}\right)$$

it is easy to verify the equivalence of problem 2.1 to the following algebraic system

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{x}{2}, -\frac{x}{2}\right) = \gamma_1(x) - \alpha_1(x) \left[ u\left(\frac{1}{2}, \frac{1}{2}\right) - u(0, 0) \right],$$

$$\alpha_2(x) u\left(\frac{x}{2}, \frac{x}{2}\right) + u\left(\frac{x}{2}, -\frac{x}{2}\right) = \gamma_2(x) - \alpha_2(x) \left[ u\left(\frac{1}{2}, -\frac{1}{2}\right) - u(0, 0) \right],$$

which by virtue of (2), (3) is unambiguously and unconditionally solvable in the class  $C(\bar{J}) \cap C^2_2(J)$ .

Therefore, problem 2.1 is reduced in an equivalent way to the Goursat problem with data on  $AB$  and  $AD$ , which is known to be correct.

*Problem 2.2.* Find a regular  $\Omega_1$  solution  $u(x, y)$  of equation (1), the domain satisfying the conditions

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{1+x}{2}, \frac{1-x}{2}\right) = \gamma_1(x), \quad x \in \bar{J},$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{x+1}{2}, \frac{x-1}{2}\right) = \gamma_2(x), \quad x \in \bar{J},$$

where

$$\alpha_1(0) \cdot \alpha_1(1) \neq \alpha_2(0) \cdot \alpha_2(1), \tag{4}$$

$$1 + \alpha_1(x) \neq 0, 1 + \alpha_2(x) \neq 0 \quad \forall x \in \bar{J} \tag{5}$$

$$\alpha_i(x), \gamma_i(x) \in C(\bar{J}) \cap C^2(J), \quad i = 1, 2.$$

Based on the Asgerisson principle for characteristic quadrilaterals with vertices respectively at points

$$\left(\frac{x}{2}, \frac{x}{2}\right), (1, 0), \left(\frac{x+1}{2}, \frac{x-1}{2}\right), \left(\frac{1}{2}, \frac{1}{2}\right)$$

and

$$\left(\frac{x}{2}, -\frac{x}{2}\right), (1, 0), \left(\frac{1+x}{2}, \frac{1-x}{2}\right), \left(\frac{1}{2}, -\frac{1}{2}\right)$$

we are convinced of the equivalence of problem 2.2 to the following two algebraic systems for finding  $u\left(\frac{x}{2}, \frac{x}{2}\right)$ ,  $u\left(\frac{x}{2}, -\frac{x}{2}\right)$ ,  $u\left(\frac{x+1}{2}, \frac{x-1}{2}\right)$  and  $u\left(\frac{1+x}{2}, \frac{1-x}{2}\right)$

$$\begin{cases} u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{x+1}{2}, \frac{x-1}{2}\right) = \gamma_1(x), \\ u\left(\frac{x}{2}, \frac{x}{2}\right) - u\left(\frac{x+1}{2}, \frac{x-1}{2}\right) = u\left(\frac{1}{2}, \frac{1}{2}\right) - u(1, 0) \end{cases}$$

and

$$\begin{cases} u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{1+x}{2}, \frac{1-x}{2}\right) = \gamma_2(x), \\ u\left(\frac{x}{2}, -\frac{x}{2}\right) - u\left(\frac{1+x}{2}, \frac{1-x}{2}\right) = u\left(\frac{1}{2}, -\frac{1}{2}\right) - u(1, 0), \end{cases}$$

which, by virtue of conditions (4), (5) are uniquely solvable.

Therefore, problem 2.2 is reduced in an equivalent way, as in the case of problem 2.1, to the Goursat problem.

*Problem 2.3.* Find a regular  $\Omega_1$  solution  $u(x, y)$  of equation (1), in the domain of equation (1), satisfying the conditions

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{1-x}{2}, -\frac{1-x}{2}\right) = \gamma_1(x), \quad x \in \bar{J}, \tag{6}$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma_2(x), \quad x \in \bar{J}, \tag{7}$$

where

$$1 - \alpha_1(1-x) \alpha_2(x) \neq 0, \quad x \in \bar{J}, \tag{8}$$

$$\alpha_i(x), \gamma_i(x) \in C(\bar{J}) \cap C^2(J), \quad i = 1, 2.$$

Replacing in (6) everywhere  $x$  by  $1-x$ , to find  $u\left(\frac{x}{2}, -\frac{x}{2}\right)$  we obtain the following algebraic system

$$\alpha_1(1-x) u\left(\frac{x}{2}, -\frac{x}{2}\right) + u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma_1(1-x),$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma_2(x).$$

Similarly, replacing in (7)  $x$  by  $1 - x$ , to find  $u\left(\frac{x}{2}, \frac{x}{2}\right)$  we get

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{1-x}{2}, -\frac{1-x}{2}\right) = \gamma_1(x),$$

$$\alpha_2(1-x) u\left(\frac{x}{2}, \frac{x}{2}\right) + u\left(\frac{1-x}{2}, -\frac{1-x}{2}\right) = \gamma_2(1-x).$$

The last two systems are unconditionally and unambiguously solvable under the conditions (8). Therefore, problem 2.3 is reduced equivalently, as in the case of problem 2.1, to the Goursat problem.

*Problem 2.4.* Find regular in the area  $\Omega_1$  solution  $u(x, y)$  of equation (1), satisfying conditions

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{2-x}{2}, \frac{x}{2}\right) = \gamma_1(x), \quad x \in \bar{J},$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{2-x}{2}, -\frac{x}{2}\right) = \gamma_2(x), \quad x \in \bar{J},$$

where  $\alpha_i(x), \gamma_i(x) \in C(\bar{J}) \cap C^2(J), i = 1, 2$ .

Based on Asgerisson's principle for characteristic quadrilaterals with vertices respectively at points

$$(0, 0), \left(\frac{1}{2}, \frac{1}{2}\right), \left(\frac{2-x}{2}, \frac{x}{2}\right), \left(\frac{1-x}{2}, -\frac{1-x}{2}\right)$$

and

$$(0, 0), \left(\frac{1}{2}, -\frac{1}{2}\right), \left(\frac{2-x}{2}, -\frac{x}{2}\right), \left(\frac{1-x}{2}, \frac{1-x}{2}\right),$$

we are convinced of the equivalence of problem 2.4 to the following algebraic system

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{1-x}{2}, -\frac{1-x}{2}\right) = \gamma_1(x) - \alpha_1(x) \left[ u\left(\frac{1}{2}, \frac{1}{2}\right) - u(0, 0) \right],$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma_2(x) - \alpha_2(x) \left[ u\left(\frac{1}{2}, -\frac{1}{2}\right) - u(0, 0) \right].$$

Therefore, problem 2.4 is equivalently reduced to problem 2.3.

*Problem 2.5.* Find regular in the area  $\Omega_1$  solution  $u(x, y)$  of equation (1), satisfying conditions

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{2-x}{2}, -\frac{x}{2}\right) = \gamma_1(x), \quad x \in \bar{J},$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{2-x}{2}, \frac{x}{2}\right) = \gamma_2(x), \quad x \in \bar{J},$$

where

$$\alpha_1(0) \neq \alpha_2(0), \alpha_1(1) \cdot \alpha_2(1) \neq 1, \tag{9}$$

$$\alpha_1(x) \cdot \alpha_1(1-x) \neq 1, \alpha_2(x) \cdot \alpha_2(1-x) \neq 1 \quad \forall x \in \bar{J}, \tag{10}$$

$$\alpha_i(x), \gamma_i(x) \in C(\bar{J}) \cap C^2(J), \quad i = 1, 2.$$

Using the same reasoning as in problem 2.4, we are convinced that problem 2.5 is equivalent to the following algebraic system

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma_1(x) - \alpha_1(x) \left[ u\left(\frac{1}{2}, -\frac{1}{2}\right) - u(0, 0) \right],$$

$$u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{1-x}{2}, \frac{x-1}{2}\right) = \gamma_2(x) - \alpha_2(x) \left[ u\left(\frac{1}{2}, \frac{1}{2}\right) - u(0, 0) \right],$$

which is apparently equivalently reduced to the next two algebraic systems

$$\begin{cases} u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha_1(x) u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma_1(x) - \alpha_1(x) \left[ u\left(\frac{1}{2}, -\frac{1}{2}\right) - u(0, 0) \right], \\ \alpha_1(1-x) u\left(\frac{x}{2}, \frac{x}{2}\right) + u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma_1(1-x) - \alpha_1(1-x) \left[ u\left(\frac{1}{2}, -\frac{1}{2}\right) - u(0, 0) \right] \end{cases}$$

and

$$\begin{cases} u\left(\frac{x}{2}, -\frac{x}{2}\right) + \alpha_2(x) u\left(\frac{1-x}{2}, \frac{x-1}{2}\right) = \gamma_2(x) - \alpha_2(x) \left[ u\left(\frac{1}{2}, \frac{1}{2}\right) - u(0, 0) \right], \\ \alpha_2(1-x) u\left(\frac{x}{2}, -\frac{x}{2}\right) + u\left(\frac{1-x}{2}, \frac{x-1}{2}\right) = \gamma_2(1-x) - \alpha_2(1-x) \left[ u\left(\frac{1}{2}, \frac{1}{2}\right) - u(0, 0) \right]. \end{cases}$$

If the conditions (9) and (10) are met, the last two systems are unambiguously and unconditionally solvable. Therefore, problem 2.5, as in the case of problem 2.1, is equivalently reduced to Goursat problem.

### 3 Non-local problems with intra-boundary displacement

*Problem 3.1.* Find regular in the area  $\Omega_1$  solution  $u(x, y)$  of equation (1), satisfying conditions

$$\begin{aligned} u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha u\left(\frac{1}{2}, \frac{1}{2} - x\right) &= \gamma(x), \quad x \in \bar{J}, \\ u(x, 0) &= \tau(x), \quad x \in \bar{J}, \end{aligned} \tag{11}$$

where

$$\alpha \neq \frac{1}{2}, \alpha [\gamma(1) - \alpha\tau(1)] = (\alpha^2 + \alpha - 1)\tau(0) - (1 - \alpha)\gamma(0), \tag{12}$$

$$\tau(x), \gamma(x) \in C(\bar{J}) \cap C^2(J). \tag{13}$$

Based on Asgeirsson's principle for characteristic quadrilaterals with vertices respectively at points

$$\left(\frac{x}{2}, \frac{x}{2}\right), \left(\frac{1}{2}, \frac{1}{2} - x\right), \left(\frac{1-x}{2}, \frac{1-x}{2}\right) \text{ and } (x, 0),$$

seeing the equivalence of problem 3.1 to the following algebraic system

$$\begin{aligned} u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha u\left(\frac{1}{2}, \frac{1}{2} - x\right) &= \gamma(x), \\ u\left(\frac{x}{2}, \frac{x}{2}\right) + u\left(\frac{1}{2}, \frac{1}{2} - x\right) &= u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) + u(x, 0). \end{aligned}$$

From where

$$(1 - \alpha) u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma(x) - \alpha\tau(x). \tag{14}$$

Changing everywhere  $x$  on  $1-x$ , from the last equation we get

$$\alpha u\left(\frac{x}{2}, \frac{x}{2}\right) + (1 - \alpha) u\left(\frac{1-x}{2}, \frac{1-x}{2}\right) = \gamma(1-x) - \alpha\tau(1-x). \tag{15}$$

From the system (14), (15) we find that

$$u\left(\frac{x}{2}, \frac{x}{2}\right) = \frac{1 - \alpha}{1 - 2\alpha} [\gamma(x) - \alpha\tau(x)] - \frac{\alpha}{1 - 2\alpha} [\gamma(1-x) - \alpha\tau(1-x)]. \tag{16}$$

Consequently, the solution of problem 3.1 is equivalently reduced to the problem of Darboux (11), (16), the regular solution of which, when the conditions (12), (13) are met, exists only. Next, obtained in the area  $\Omega_2$  the solution can be naturally continued throughout  $\Omega_1$ .

*Problem 3.2.* Find regular in the area  $\Omega_2$  solution  $u(x, y)$  of equation (1), satisfying conditions

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \alpha u\left(\frac{1}{2}, \left|\frac{1}{2} - x\right|\right) = \gamma(x), \quad x \in \bar{J},$$

$$u(x, 0) = \tau(x), \quad x \in \bar{J}.$$

By virtue of the Asgeirsson principle  $0 < x^* < \frac{1}{2}$  fair ratio

$$u\left(\frac{x^*}{2}, \frac{x^*}{2}\right) + u\left(\frac{1}{2}, \left|\frac{1}{2} - x^*\right|\right) = u(x^*, 0) + u\left(\frac{1-x^*}{2}, \frac{1-x^*}{2}\right).$$

For a symmetric point with respect to zero, the following ratio is valid

$$u\left(\frac{x^*}{2}, \frac{x^*}{2}\right) + u\left(\frac{1}{2}, \left|\frac{1}{2} - x^*\right|\right) = u(1-x^*, 0) + u\left(\frac{1-x^*}{2}, \frac{1-x^*}{2}\right).$$

This suggests that the mean theorem, which is valid for all  $x \in \bar{J}$  we can only use it when

$$u(x, 0) = u(1-x, 0).$$

*Theorem 3.1.* Let  $\tau(x), \gamma(x) \in C(\bar{J}) \cap C^2(J)$  and the conditions are met

$$\alpha \neq 1, \tau(x) \equiv \tau(1-x) \quad \forall x \in \bar{J}. \tag{17}$$

Then there is only one regular in the field  $\Omega_2$  problem solving 3.2.

When the conditions (17) are met, it is sufficient to repeat the same reasoning and calculations as in the study of problem 3.1 to prove the theorem 3.1.

*Problem 3.3.* Find a regular solution in the region of  $\Omega_2$  of the  $u(x, y)$  equation (1) satisfying the conditions

$$u\left(\frac{x}{2}, \frac{x}{2}\right) + \beta u\left(\frac{1}{2}, \frac{x}{2}\right) = \gamma(x), \quad x \in \bar{J}, \tag{18}$$

$$u(x, 0) = \tau(x), \quad x \in \bar{J}. \tag{19}$$

*Theorem 3.2.* Let  $|\beta| < 1, \tau(x), \gamma(x) \in C(\bar{J}) \cap C^2(J)$  and the condition is fulfilled

$$\tau(0) + \beta\tau\left(\frac{1}{2}\right) = \gamma(0).$$

Then problem 3.3 is uniquely solvable, its solution is represented as

$$u(x, y) = \tau(x-y) + \beta \left[ \tau\left(\frac{1-x+y}{2}\right) - \tau\left(\frac{1-x-y}{2}\right) \right] - \gamma(x-y) + \gamma(x+y) -$$

$$- \sum_{j=1}^{\infty} \sum_{i=j}^{\infty} (-1)^i \beta^i \left[ \gamma\left(\frac{(2j-1)+x-y}{2^i}\right) - \gamma\left(\frac{(2j-1)+x+y}{2^i}\right) - \right.$$

$$\left. - \gamma\left(\frac{(2j-1)-x+y}{2^i}\right) + \gamma\left(\frac{(2j-1)-x-y}{2^i}\right) + \right. \tag{20}$$

$$+ \beta\tau\left(\frac{2(2j-1)-1+x-y}{2^{i+1}}\right) - \beta\tau\left(\frac{2(2j-1)+1+x+y}{2^{i+1}}\right) -$$

$$\left. - \beta\tau\left(\frac{2(2j-1)-1-x+y}{2^{i+1}}\right) + \beta\tau\left(\frac{2(2j-1)+1-x-y}{2^{i+1}}\right) \right].$$

*Proof.* It is known that any regular solution of the equation (1) can be represented as

$$u(x, y) = f(x - y) + g(x + y), \quad (21)$$

where  $f(x), g(x) \in C^2(\Omega) \cap C(\bar{\Omega})$ .

Satisfying (21) conditions (18), (19) for finding  $f(x)$  and  $g(x)$  we obtain the following system of functional equations

$$f(x) = \tau(x) - g(x), \quad (22)$$

$$g(x) - \beta g\left(\frac{1-x}{2}\right) + \beta g\left(\frac{1+x}{2}\right) = \gamma(x) - \beta \tau\left(\frac{1-x}{2}\right). \quad (23)$$

The solution of equation (23) under the condition  $|\beta| < 1$  can be constructed by the iteration method. Indeed

$$g_0(x) = \gamma(x) - \beta \tau\left(\frac{1-x}{2}\right),$$

$$g_1(x) = \gamma(x) - \beta \tau\left(\frac{1-x}{2}\right) + \beta \gamma\left(\frac{1-x}{2}\right) - \beta \gamma\left(\frac{1+x}{2}\right) - \\ - \beta^2 \tau\left(\frac{1+x}{4}\right) + \beta^2 \tau\left(\frac{3-x}{4}\right)$$

etc.

Continuing this process indefinitely, we get

$$g(x) = \gamma(x) - \beta \tau\left(\frac{1-x}{2}\right) + \\ + \sum_{j=1}^{\infty} \sum_{i=j}^{\infty} (-1)^i \beta^i \left[ \gamma\left(\frac{(2j-1)+x}{2^i}\right) - \gamma\left(\frac{(2j-1)-x}{2^i}\right) + \right. \\ \left. + \beta \tau\left(\frac{2(2j-1)-1+x}{2^{i+1}}\right) - \beta \tau\left(\frac{2(2j-1)+1-x}{2^{i+1}}\right) \right]. \quad (24)$$

It is easily verified that the series itself (24) the series obtained after differentiation converge uniformly. Substituting (24) into (22) we find  $f(x)$ , and hence  $u(x, y)$  by formula (21).

It should be noted that formula (20) in the case of  $\beta = 0$  coincides with the formula for solving the Darboux problem.

### Conclusion

In this paper, all problems are considered in the characteristic quadrilateral; therefore, the choice of an inner manifold, the points of which are associated with the boundary manifold, is small. In the case of sufficiently derived domains (for example, a rectangular noncharacteristic domain), a problem arises both in the nonlocal condition itself and in the corresponding inner manifold. This method is of particular interest for weakly hyperbolic equations.

### References

- 1 Нахушев А.М. О некоторых краевых задачах для гиперболических уравнений и уравнений смешанного типа / А.М. Нахушев // Дифференц. уравнения. — Т. 5. — № 1. — 1969. — С. 44–59.

- 2 Нахушев А.М. Уравнения математической биологии / А.М. Нахушев. — М.: Высш. шк., 1985. — 301 с.
- 3 Нахушев А.М. Задачи со смещением для уравнений в частных производных / А.М. Нахушев. — М.: Наука, 2002. — 287 с.
- 4 Кальменов Т.Ш. К теории начально-краевых задач для дифференциальных уравнений / Т.Ш. Кальменов. — Алматы: Ин-т мат. и мат. моделир., 2013. — 406 с.
- 5 Нахушев А.М. Методика постановки корректных краевых задач для линейных гиперболических уравнений второго порядка на плоскости / А.М. Нахушев // Дифференц. уравнения. — Т. 6. — № 1. — 1970. — С. 192–195.
- 6 Нахушев А.М. О задаче Дарбу для гиперболических уравнений / А.М. Нахушев // ДАН СССР. — Т. 195. — № 4. — 1970. — С. 776–779.
- 7 Кальменов Т.Ш. О регулярных краевых задачах для волнового уравнения / Т.Ш. Кальменов // Дифференц. уравнения. — Т. 17. — № 6. — 1981. — С. 1105–1121.
- 8 Кальменов Т.Ш. Спектр краевой задачи со смещением для волнового уравнения / Т.Ш. Кальменов // Дифференц. уравнения. — Т. 19. — № 1. — 1983. — С. 75–78.
- 9 Аттаев А.Х. Краевые задачи с внутренне-краевым смещением для уравнения колебания струны / А.Х. Аттаев // Докл. Адыгской (Черкесской) Междунар. акад. наук. — Т. 16. — № 2. — 2014. — С. 17–19.
- 10 Дезин А.А. К общей теории граничных задач / А.А. Дезин // Мат. сб. — Т. 100(142). — № 2(6). — 1976. — С. 171–180.
- 11 Дезин А.А. Об операторных уравнениях второго порядка / А.А. Дезин // Сиб. мат. журн. — Т. 19. — № 5. — 1978. — С. 1032–1042.

А.Х. Аттаев

*РҒА Кабардин-Балқар ғылыми орталығының Қолданбалы математика және автоматтандыру институты,  
Нальчик, Ресей*

## Ішек тербелісінің теңдеуі үшін кейбір бейлокал шеттік және ішкі шеттік есептер

Жұмыста классикалық Гурса және Дарбу есептерін жалпылайтын бірқатар бейлокал есептер тұжырымдалған және зерттелген. Олардың кейбіреулері шеттік, ал екінші бөлігі ішкі шеттік болып табылады және екі жағдайда да сипаттамалық және сипаттамалық емес ығысулар қарастырылған. Сондай-ақ, мақалада қарастырылған бірқатар есептер ішек тербелісінің модельдік жүктелген теңдеуі үшін корректілі қойылған есептер теориясын құруда ерекше жағдай ретінде туындағанын атап өткен жөн.

*Кілт сөздер:* толқындық теңдеу, жалпы шешім, Коши есебі, Гурса есебі, Дарбу есебі, сипаттамалық ығысу есептері, сипаттамалық емес ығысу есептері.



А.Х. Аттаев

*Институт прикладной математики и автоматизации Кабардино-Балкарского научного центра РАН,  
Нальчик, Россия*

## О некоторых нелокальных краевых и внутренних краевых задачах для уравнения колебания струны

В работе сформулированы и исследованы некоторые нелокальные задачи, обобщающие классические задачи Гурса и Дарбу. Часть из них являются краевыми, а другая — внутренними краевыми, причем в обоих случаях рассмотрены как характеристические, так и нехарактеристические смещения. Следует также отметить, что ряд задач, рассмотренных в статье, возникли как частный случай при построении теории корректных задач для модельного нагруженного уравнения колебания струны.

*Ключевые слова:* волновое уравнение, общее решение, задача Коши, задача Гурса, задача Дарбу, задача с характеристическим смещением, задача с нехарактеристическим смещением.

### References

- 1 Nakhushev, A.M. (1969). O nekotorykh kraevykh zadachakh dlia giperbolicheskikh uravnenii i uravnenii smeshannogo tipa [On some boundary value problems for hyperbolic equations and mixed-type equations]. *Differentsialnye uravneniia — Differential Equations*, 5(1), 44–59 [in Russian].
- 2 Nakhushev, A.M. (1985). *Uravneniia matematicheskoi biologii [Equations of mathematical biology]*. Moscow: Vysshiaia shkola [in Russian].
- 3 Nakhushev, A.M. (2002). *Zadachi so smeshcheniem dlia uravnenii v chastnykh proizvodnykh [Displacement problems for partial differential equations]*. Moscow: Nauka [in Russian].
- 4 Kalmenov, T.S. (2013). *K teorii nachalno-kraevykh zadach dlia differentsialnykh uravnenii [On the theory of initial boundary value problems for differential equations]*. Almaty: Institut matematiki i matematicheskogo modelirovaniia [in Russian].
- 5 Nakhushev, A.M. (1970). Metodika postanovki korrektnykh kraevykh zadach dlia lineinykh giperbolicheskikh uravnenii vtorogo poriadka na ploskosti [Methodology for setting correct boundary value problems for linear hyperbolic equations of the second order on the plane]. *Differentsialnye uravneniia — Differential Equations*, 6(1), 192–195 [in Russian].
- 6 Nakhushev, A.M. (1970). O zadache Darbu dlia giperbolicheskikh uravnenii [On the Darboux problem for hyperbolic equations]. *DAN SSSR — DAN USSR*, 195(4), 776–779 [in Russian].
- 7 Kalmenov, T.S. O reguliarnykh kraevykh zadachakh dlia volnovogo uravneniia [On regular boundary value problems for the wave equation]. *Differentsialnye uravneniia — Differential Equations*, 17(6), 1105–1121 [in Russian].
- 8 Kalmenov, T.S. (1983). Spektr kraevoi zadachi so smeshcheniem dlia volnovogo uravneniia [Spectrum of the boundary value problem with displacement for the wave equation]. *Differentsialnye uravneniia — Differential Equations*, 19(1), 75–78 [in Russian].
- 9 Attaev, A.H. (2014). Kraevye zadachi s vnutrenne-kraevym smeshcheniem dlia uravneniia kolebaniia struny [Boundary value problems with internal-boundary displacement for the string oscillation equation]. *Doklady Adygskoi (Cherkesskoi) Mezhdunarodnoi akademii nauk — Reports of the Adyghe (Circassian) International Academy of Sciences*, 16(2), 17–19 [in Russian].
- 10 Dezin, A.A. (1976). K obshchei teorii granichnykh zadach [To the general theory of boundary value problems]. *Matematicheskii sbornik — Mathematical collection*, 100(142), 2 (6), 171–180 [in Russian].

- 11 Dezin, A.A. (1978). Ob operatornykh uravneniakh vtorogo poriadka [Second-order operator equations]. *Sibirskii matematicheskii zhurnal — Siberian Mathematical Journal*, 19(5), 1032–1042 [in Russian].