

On the formulation and investigation of a boundary value problem for a third-order equation of a parabolic-hyperbolic type

M. Mamajonov¹, Q. Rakhimov^{2,*}, Kh. Shermatova²

¹Kokand State Pedagogical Institute, Kokand, Uzbekistan;

²Fergana State University, Fergana, Uzbekistan

(E-mail: mirzamamajonov@gmail.com, quvvatali.rahimov@gmail.com, shermatovahilola1978@gmail.com)

In the paper a novel boundary value problem for a third-order partial differential equation (PDE) of a parabolic-hyperbolic type, within a pentagonal domain consisting of both parabolic and hyperbolic regions was investigated. Such equations are pivotal in modeling complex physical phenomena across diverse fields such as physics, engineering, and finance due to their ability to encapsulate a wide range of dynamics through their mixed-type nature. By employing a constructive solution approach, we demonstrate the unique solvability of the posed problem. The significance of this study lies in its extension of the mathematical framework for understanding and solving higher-order mixed PDEs in complex geometrical domains, thus offering new avenues for theoretical and applied research in mathematical physics and related disciplines.

Keywords: differential equations, parabolic-hyperbolic type, a third-order parabolic-hyperbolic type.

2020 Mathematics Subject Classification: 35K30, 35L20, 35R05, 35J25.

Introduction

The study of non-classical equations of mathematical physics refers to the investigation of partial differential equations (PDEs) that exhibit behaviors beyond the standard classifications of parabolic, hyperbolic, and elliptic equations. These equations are often referred to as non-classical or degenerate equations. At present, the study of non-classical equations of mathematical physics is being intensively developed — equations of mixed, composite and mixed-composite types. One of the main reasons is the emergence of applied applications of boundary value problems posed for equations of these types. Many problems in physics, technology, mechanics and other areas require the study of such equations.

First, they began to study second-order mixed equations of the elliptic-hyperbolic type. The Italian mathematician Tricomi began to study fundamental studies of equations of such types in the 1920s.

After that, we began to study many different problems for equations of these types. A review of theoretical and applied research is given in the works and books of A.V. Bitsadze, L. Bers, M.M. Smirnov, as well as, in Uzbekistan, in the books of M.S. Salokhitdinov, T.D. Juraev.

Research into equations of elliptic-parabolic, parabolic-hyperbolic types began in the 1950s and 1960s. In 1959, I.M. Gelfand [1] pointed out the need for joint consideration of equations in one part of the domain of parabolic, and the other part of hyperbolic types. He gives an example related to the movement of gas in a channel surrounded by a porous medium: in the channel, the movement of gas is described by the wave equation, and outside it — by the diffusion equation.

Then, in the 1970s and 1980s, they began to study various problems for equations of the third and higher orders of the parabolic-hyperbolic type. Such problems were studied mainly by T.D. Dzhuraev and his students (for example, see [2, 3]).

At present, the study of various boundary value problems for equations of the third and higher orders of the parabolic-hyperbolic type has been developed on a broad scale (for example, see [4–15]).

*Corresponding author. E-mail: quvvatali.rahimov@gmail.com

Received: 05 January 2024; Accepted: 04 March 2024.

© 2024 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1 Formulation of the problem

In this article, we consider one boundary value problem for a third-order parabolic-hyperbolic type equation of the form

$$\left(\frac{\partial}{\partial x} - \frac{\partial}{\partial y} + c\right)(Lu) = 0 \tag{1}$$

in the pentagonal region G of the plane xOy , where $G = G_1 \cup G_2 \cup G_3 \cup J_1 \cup J_2$,

$$Lu = \begin{cases} u_{xx} - u_y, & (x, y) \in D_1, \\ u_{xx} - u_{yy}, & (x, y) \in D_i, \quad i = 2, 3, \end{cases}$$

$c \in R$, and G_1 is a rectangle with vertices at points $A(0;0)$, $B(1;0)$, $B_0(1,1)$, $A_0(0,1)$; G_2 — triangle with vertices at points B , $C(0,-1)$, $D(-1,0)$; G_3 — rectangle with vertices at points A , D , $D_0(-1,1)$, A_0 ; J_1 — open segment with vertices at points B , D ; J_2 — an open segment with vertices at points A , A_0 .

The equation (1) is a special case of the equation $\left(a\frac{\partial}{\partial x} + b\frac{\partial}{\partial y} + c\right)(Lu) = 0$ when $\gamma = \frac{b}{a} = -1$, that is, the angular coefficient of the characteristic of the operator $a\frac{\partial}{\partial x} + b\frac{\partial}{\partial y}$ is equal to $\gamma = -1$.

For the equation (1), the following problem is posed:

Problem 1. It is required to find the function $u(x, y)$ which is 1) continuous in \overline{G} and in the domain of $G \setminus J_1 \setminus J_2$ has continuous derivatives involved in the equation (1), and u_x and u_y are continuous in G up to part of the boundary of the domain G specified in the boundary conditions; 2) satisfies the equation (1) in the domain $G \setminus J_1 \setminus J_2$; 3) satisfies the following boundary conditions:

$$u(1, y) = \varphi_1(y), \quad 0 \leq y \leq 1, \tag{2}$$

$$u(-1, y) = \varphi_2(y), \quad 0 \leq y \leq 1,$$

$$u_x(1, y) = \varphi_3(y), \quad 0 \leq y \leq 1,$$

$$u|_{BC} = \psi_1(x), \quad 0 \leq x \leq 1, \tag{3}$$

$$u|_{DF} = \psi_2(x), \quad -1 \leq x \leq -\frac{1}{2}, \tag{4}$$

$$\left.\frac{\partial u}{\partial n}\right|_{BC} = \psi_3(x), \quad -1 \leq x \leq 0; \tag{5}$$

4) satisfies the following gluing conditions on the lines of type changing:

$$u(x, +0) = u(x, -0) = T(x), \quad -1 \leq x \leq 1, \tag{6}$$

$$u_y(x, +0) = u_y(x, -0) = N(x), \quad -1 \leq x \leq 1, \tag{7}$$

$$u_{yy}(x, +0) = u_{yy}(x, -0) = M(x), \quad -1 \leq x \leq 1, \tag{8}$$

$$u(+0, y) = u(-0, y) = \tau_3(y), \quad 0 \leq y \leq 1, \tag{9}$$

$$u_x(+0, y) = u_x(-0, y) = \nu_3(y), \quad 0 \leq y \leq 1, \tag{10}$$

$$u_{xx}(+0, y) = u_{xx}(-0, y) = \mu_3(y), \quad 0 \leq y \leq 1, \tag{11}$$

where

$$T(x) = \begin{cases} \tau_1(x), & \text{if } 0 \leq x \leq 1, \\ \tau_2(x), & \text{if } -1 \leq x \leq 0; \end{cases}$$

$N(x) = \begin{cases} \nu_1(x), & \text{if } 0 \leq x \leq 1, \\ \nu_2(x), & \text{if } -1 \leq x \leq 0; \end{cases}$ $M(x) = \begin{cases} \mu_1(x), & \text{if } 0 < x < 1, \\ \mu_2(x), & \text{if } -1 < x < 0, \end{cases}$ φ_i, ψ_i ($i = 1, 2, 3$) are given sufficiently smooth functions, τ_i, ν_i, μ_i ($i = 1, 2, 3$) are unknown yet sufficiently smooth functions, n is an internal normal to the line $x - y = 1$, and the point F has coordinates $F(-1/2, -1/2)$.

2 Studying of the Problem

Theorem 2.1. If $\varphi_1, \varphi_2 \in C^3[0, 1]$, $\varphi_3 \in C^2[0, 1]$, $\psi_1 \in C^3[0, 1]$, $\psi_2 \in C^3[-1, -1/2]$, $\psi_3 \in C^2[0, 1]$, and the matching conditions $\varphi_1(0) = \psi_1(1)$ fulfilled, $\psi_2(-1) = \varphi_2(0)$, then Problem 1 is uniquely solvable.

Proof. We will prove the theorem by constructing the solution. To do this, we will rewrite the equation (1) as

$$u_{1xx} - u_{1y} = \omega_1(x + y) \exp(cy), \quad (x, y) \in G_1, \tag{12}$$

$$u_{ixx} - u_{iyy} = \omega_i(x + y) \exp(cy), \quad (x, y) \in G_i, \quad i = 2, 3, \tag{13}$$

where the notation $u(x, y) = u_i(x, y)$, $(x, y) \in G_i$ ($i = \overline{1, 3}$), functions $\omega_i(x + y)$, $i = \overline{1, 3}$ are unknown sufficiently smooth functions to be determined.

The study will be carried out first in the domain G_2 . The solution of the equation (13) ($i = 2$) satisfying the conditions (6), (7) can be represented as

$$u_2(x, y) = \frac{1}{2}[T(x + y) + T(x - y)] + \frac{1}{2} \int_{x-y}^{x+y} N(t) dt - \frac{1}{2} \int_0^y \exp(c\eta) d\eta \int_{x-y+\eta}^{x+y-\eta} \omega_2(\xi + \eta) d\xi. \tag{14}$$

Substituting (14) into the condition (5), after some calculations, we find

$$\omega_2(x + y) = -\sqrt{2}\psi'_3\left(\frac{x + y + 1}{2}\right) \exp\left[-\frac{c}{2}(x + y - 1)\right], \quad -1 \leq x + y \leq 1.$$

Further, substituting (14) into the condition (3), after some simplifications, we obtain the first relation between the unknown functions $T(x)$ and $N(x)$ on the line J_1 of type changing:

$$T'(x) + N(x) = \alpha_1(x), \quad -1 \leq x \leq 1, \tag{15}$$

where $\alpha_1(x) = \psi'_1\left(\frac{x+1}{2}\right) + \omega_2(x) \int_0^{(x-1)/2} e^{c\eta} d\eta$.

If we take into account the representation of the function $T(x)$, then for $-1 \leq x \leq 0$ the equation (15) has the form

$$\tau'_2(x) + \nu_2(x) = \alpha_1(x), \quad -1 \leq x \leq 0. \tag{16}$$

Now, substituting (14) into (4), after some transformations, we have

$$\tau'_2(x) - \nu_2(x) = \delta_1(x), \quad -1 \leq x \leq 0, \tag{17}$$

where $\delta_1(x) = \psi'_2\left(\frac{x-1}{2}\right) + \int_0^{-(x+1)/2} e^{c\eta} \omega_2(x + 2\eta) d\eta$.

From (16) and (17) we obtain

$$\tau_2(x) = \frac{1}{2} \int_{-1}^x [\alpha_1(t) + \delta_1(t)] dt + \psi_2(-1), \quad \nu_2(x) = \frac{1}{2} [\alpha_1(x) - \delta_1(x)].$$

For $0 \leq x \leq 1$ from (15), we have the first relation between the unknown functions $\tau_1(x)$ and $\nu_1(x)$ on the line J_1 of type changing in the following form:

$$\tau'_1(x) + \nu_1(x) = \alpha_1(x), \quad 0 \leq x \leq 1. \tag{18}$$

Passing to the limit at $y \rightarrow 0$ in the equation (13) ($i = 2$), we will find the second relation between the unknown functions $\tau_1(x)$ and $\mu_1(x)$ on J_1 :

$$\tau''_1(x) - \mu_1(x) = \omega_2(x), \quad 0 \leq x \leq 1. \tag{19}$$

The equation (1) in the domain G_1 can be rewritten as

$$u_{1xxx} - u_{1xy} - u_{1xxy} + u_{1yy} + cu_{1xx} - cu_{1y} = 0.$$

Passing to the limit at $y \rightarrow 0$ in the last equation, we obtain the third relation between the unknown functions $\tau_1(x)$, $\nu_1(x)$ and $\mu_1(x)$ on the line of type changing J_1 :

$$\tau_1'''(x) - \nu_1'(x) - \nu_1''(x) + \mu_1(x) + c\tau_1''(x) - c\nu_1(x) = 0, \quad 0 \leq x \leq 1. \quad (20)$$

Eliminating the functions $\nu_1(x)$ and $\mu_1(x)$ from the equations (18), (19) and (20) and integrating the resulting equation from 0 to x , we arrive at the equation

$$\tau_1''(x) + (1 + \frac{c}{2})\tau_1'(x) + \frac{c}{2}\tau_1(x) = \alpha_2(x) + k_1, \quad 0 \leq x \leq 1, \quad (21)$$

where $\alpha_2(x) = \frac{1}{2}\alpha_1'(x) + \frac{1}{2}\alpha_1(x) + \frac{1}{2}\int_0^x [\omega_2(t) + c\alpha_1(t)] dt$, and k_1 is still unknown constant.

When solving the equation (21), we consider the following cases:

- 1°. $c \neq 2, c \neq 0$;
- 2°. $c = 2$;
- 3°. $c = 0$.

In the case 1° it is easy to see that the solution of the equation (21) satisfying the conditions

$$\begin{aligned} \tau_1(0) &= \frac{1}{2} \int_{-1}^0 [\alpha_1(t) + \delta_1(t)] dt + \psi_2(-1), \\ \tau_1'(0) &= \frac{1}{2} [\alpha_1(0) + \delta_1(0)], \\ \tau_1(1) &= \varphi_1(0) \end{aligned} \quad (22)$$

has the form

$$\begin{aligned} \tau_1(x) &= \frac{2}{2-c} \int_0^x [e^{\frac{c}{2}(t-x)} - e^{t-x}] \alpha_2(t) dt + \\ &+ \frac{2k_1}{2-c} \left[\frac{2}{c} (1 - e^{-\frac{c}{2}x}) - (1 - e^{-x}) \right] + k_2 e^{-x} + k_3 e^{-\frac{c}{2}x}, \end{aligned}$$

where $k_3 = \frac{1}{2-c} \left\{ \int_{-1}^0 [\alpha_1(t) + \delta_1(t)] dt + 2\psi_2(-1) + \alpha_1(0) + \delta_1(0) \right\}$,

$$k_2 = \frac{1}{c-2} \left\{ \frac{c}{2} \int_{-1}^0 [\alpha_1(t) + \delta_1(t)] dt + c\psi_2(-1) + \alpha_1(0) + \delta_1(0) \right\},$$

$$\begin{aligned} k_1 &= \left[\frac{c}{2} (1 - e^{-\frac{c}{2}}) - (1 - e^{-1}) \right]^{-1} \left\{ \frac{2-c}{2} [\varphi_1(0) - \right. \\ &\left. - k_2 e^{-1} + k_3 e^{-\frac{c}{2}}] - \int_0^1 [e^{\frac{c}{2}(t-1)} - e^{t-1}] \alpha_2(t) dt \right\}. \end{aligned}$$

Also, in the case 2°, one can show that the solution of solving the equation (21) satisfying the conditions (22), has the following form

$$\tau_1(x) = \int_0^x (x-t)e^{t-x} \alpha_2(t) dt + k_1 [1 - (x+1)e^{-x}] + (k_2 + k_3 x)e^{-x},$$

where $k_2 = \frac{1}{2} \int_{-1}^0 [\alpha_1(t) + \delta_1(t)] dt + \psi_2(-1)$, $k_3 = k_2 + \frac{1}{2} [\alpha_1(0) + \delta_1(0)]$,

$$k_1 = \frac{1}{e-2} \left[\varphi_1(0)e - k_2 - k_3 - \int_0^1 (1-t)e^t \alpha_2(t) dt \right].$$

Moreover, for the case 3°, the solution of (21) satisfying (22) defined by

$$\tau_1(x) = \int_0^x e^{t-x} \alpha_3(t) dt + k_1(x - 1 + e^{-x}) + k_2(1 - e^{-x}) + k_3 e^{-x},$$

where $\alpha_3(x) = \int_0^x \alpha_2(t) dt$, $k_3 = \frac{1}{2} \int_{-1}^0 [\alpha_1(t) + \delta_1(t)] dt + \psi_2(-1)$, $k_2 = k_3 + \frac{1}{2} [\alpha_1(0) + \delta_1(0)]$,

$$k_1 = \varphi_1(0)e - k_2(e - 1) - k_3 - \int_0^1 e^t \alpha_3(t) dt.$$

Now, we consider the G_3 . Let us introduce the notation:

$$\omega_3(x + y) = \begin{cases} \omega_{31}(x + y), & -1 \leq x + y \leq 0, \\ \omega_{32}(x + y), & 0 \leq x + y \leq 1. \end{cases}$$

Then, passing to the limit at $y \rightarrow 0$, in the equations (13) ($i = 2$) and (13) ($i = 3$) due to (6)–(8), we get

$$\omega_{31}(x) = \omega_2(x), \quad -1 \leq x \leq 0.$$

Now, we consider the following problem:

$$\begin{cases} u_{3xx} - u_{3yy} = \omega_3(x + y)e^{cy}, \\ u_3(x, 0) = \tau_2(x), \quad u_{3y}(x, 0) = \nu_2(x), \quad -1 \leq x \leq 0, \\ u_3(-1, y) = \varphi_2(y), \quad u_3(0, y) = \tau_3(y), \quad 0 \leq y \leq 1. \end{cases}$$

The solution to this problem will be sought in the form

$$u_3(x, y) = u_{31}(x, y) + u_{32}(x, y) + u_{33}(x, y), \tag{23}$$

where $u_{31}(x, y)$ is the solution of the problem

$$\begin{cases} u_{31xx} - u_{31yy} = 0, \\ u_{31}(x, 0) = \tau_2(x), \quad u_{31y}(x, 0) = 0, \quad -1 \leq x \leq 0, \\ u_{31}(-1, y) = \varphi_2(y), \quad u_{31}(0, y) = \tau_3(y), \quad 0 \leq y \leq 1; \end{cases} \tag{24}$$

$u_{32}(x, y)$ is the solution of the problem

$$\begin{cases} u_{32xx} - u_{32yy} = 0, \\ u_{32}(x, 0) = 0, \quad u_{32y}(x, 0) = \nu_2(x), \quad -1 \leq x \leq 0, \\ u_{32}(-1, y) = 0, \quad u_{32}(0, y) = 0, \quad 0 \leq y \leq 1; \end{cases} \tag{25}$$

$u_{33}(x, y)$ is the solution of the problem

$$\begin{cases} u_{33xx} - u_{33yy} = \omega_3(x + y)e^{cy}, \\ u_{33}(x, 0) = 0, \quad u_{33y}(x, 0) = 0, \quad -1 \leq x \leq 0, \\ u_{33}(-1, y) = 0, \quad u_{33}(0, y) = 0, \quad 0 \leq y \leq 1. \end{cases} \tag{26}$$

Using the continuation method, we find solutions to the problems (24)–(26). The solutions can be represented as follows

$$u_{31}(x, y) = \frac{1}{2} [T_2(x + y) + T_2(x - y)], \tag{27}$$

where $T_2(x) = \begin{cases} 2\varphi_2(-1-x) - \tau_2(-2-x), & -2 \leq x \leq -1, \\ \tau_2(x), & -1 \leq x \leq 0, \\ 2\tau_3(x) - \tau_2(-x), & 0 \leq x \leq 1; \end{cases}$

$$u_{32}(x, y) = \frac{1}{2} \int_{x-y}^{x+y} N_2(t) dt, \tag{28}$$

where $N_2(x) = \begin{cases} -\nu_2(-2-x), & -2 \leq x \leq -1, \\ \nu_2(x), & -1 \leq x \leq 0, \\ -\nu_2(-x), & 0 \leq x \leq 1; \end{cases}$

$$u_{33}(x, y) = -\frac{1}{2} \int_0^y e^{c\eta} d\eta \int_{x-y+\eta}^{x+y-\eta} \Omega_3(\xi + \eta) d\xi. \tag{29}$$

Using the condition $u_{33}(-1, y) = 0$, after some transformations, from (29), we obtain

$$\frac{1}{2} \int_{-1-y}^{y-1} e^{\frac{c}{2}(y+1+z)} \Omega_3(z) dz = -\omega_{31}(y-1) \int_0^y e^{c\eta} d\eta. \tag{30}$$

Hence, by differentiating (30), we find

$$\begin{aligned} \Omega_{31}(-1-y) &= c\omega_{31}(y-1) \int_0^y e^{c\eta} d\eta - \\ &- 2\omega'_{31}(y-1) \int_0^y e^{c\eta} d\eta - 3\omega_{31}(y-1)e^{cy}. \end{aligned}$$

Now, using condition $u_{33}(0, y) = 0$, after some transformations, from (30), we have

$$\omega_{32}(y) \int_0^y e^{c\eta} d\eta = - \int_{-y}^y e^{\frac{c}{2}(y+z)} \Omega_3(z) dz.$$

Substituting (27), (28) and (29) into (23), we get

$$\begin{aligned} u_3(x, y) &= \frac{1}{2} [T_2(x+y) + T_2(x-y)] + \\ &+ \frac{1}{2} \int_{x-y}^{x+y} N_2(t) dt - \frac{1}{2} \int_0^y e^{c\eta} d\eta \int_{x-y+\eta}^{x+y-\eta} \Omega_3(\xi + \eta) d\xi. \end{aligned}$$

Differentiating this solution with respect to x and tending x to zero, and also taking into account the condition (10), after some transformations, we have the following relation:

$$\nu_3(y) = \tau'_3(y) + \tau'_2(-y) - \nu_2(-y) + \frac{1}{2} \int_{-y}^y e^{\frac{c}{2}(y+z)} \Omega_3(z) dz. \tag{31}$$

Passing to the limit at $x \rightarrow 0$ in the equations (12) and (13) ($i = 2$) and taking into account (9) and (11), we obtain

$$\mu_3(y) - \tau'_3(y) = \omega_{11}(y) \exp(-cy), \quad \mu_3(y) - \tau''_3(y) = \omega_{32}(y) e^{cy}.$$

Eliminating the function $\mu_3(y)$ from these equations, we find

$$\omega_{32}(y) = \omega_{11}(y) - [\tau''_3(y) - \tau'_3(y)] e^{-cy}. \tag{32}$$

Now, passing to the limit at $y \rightarrow 0$ in the equation (13) and taking (6) and (7) into account after replacing x with $x + y$, we obtain

$$\omega_{11}(x+y) = \tau''_1(x+y) - \nu_1(x+y), \quad 0 \leq x+y \leq 1, \tag{33}$$

where $\omega_1(x + y) = \begin{cases} \omega_{11}(x + y), & 0 \leq x + y \leq 1, \\ \omega_{12}(x + y), & 1 \leq x + y \leq 2. \end{cases}$

Finally, by substituting (33) into (32), we arrive at the relation

$$\omega_{32}(y) = \tau_1''(y) - \nu_1(y) - [\tau_3''(y) - \tau_3'(y)] e^{-cy}, \tag{34}$$

and substituting (34) into (31), after some calculations, we get

$$\nu_3(y) = \frac{1}{2}\tau_3'(y) - \frac{c-2}{4} \int_0^y e^{\frac{c}{2}(y-z)} \tau_3'(z) dz + \beta_1(y),$$

where

$$\begin{aligned} \beta_1(y) &= \tau_2'(-y) - \nu_2(-y) + \frac{1}{2}\nu_1(0)e^{\frac{c}{2}y} + \\ &+ \frac{1}{2} \int_{-y}^0 e^{\frac{c}{2}(y+z)} \omega_{31}(z) dz + \frac{1}{2} \int_0^y e^{\frac{c}{2}(y+z)} [\tau_1''(z) - \nu_1(z)] dz. \end{aligned}$$

Now, we consider the domain G_1 . The solution of the equation (12) satisfying the conditions (2), (6), (9) has the form

$$\begin{aligned} u_1(x, y) &= \left[\int_0^y \tau_3(\eta) G_\xi(x, y; \cdot, 0, \eta) d\eta - \right. \\ &- \int_0^y \varphi_1(\eta) G_\xi(x, y; 1, \eta) d\eta + \int_0^1 \tau_1(\xi) G(x, y; \xi, 0) d\xi - \\ &- \int_0^y e^{c\eta} d\eta \int_0^{1-\eta} \omega_{11}(\xi + \eta) G(x, y; \xi, \eta) d\xi - \\ &- \left. \int_0^y e^{c\eta} d\eta \int_{1-\eta}^1 \omega_{12}(\xi + \eta) G(x, y; \xi, \eta) d\xi. \right] \end{aligned}$$

Differentiating this solution by x and passing x to zero and to one, we obtain the following relations

$$\begin{aligned} \nu_3(y) &= - \int_0^y \tau_3'(\eta) N(0, y; 0, \eta) d\eta + \\ &+ \int_0^y \varphi_1'(\eta) N(0, y; 1, \eta) d\eta + \int_0^1 \tau_1'(\xi) N(0, y; \xi, 0) d\xi + \\ &+ \int_0^y e^{c\eta} d\eta \int_0^{1-\eta} [\tau_1''(\xi + \eta) - \nu_1(\xi + \eta)] N_\xi(0, y; \xi, \eta) d\xi + \\ &+ \int_0^y e^{c\eta} d\eta \int_{1-\eta}^1 \omega_{12}(\xi + \eta) N_\xi(0, y; \xi, \eta) d\xi, \\ \varphi_3(y) &= - \int_0^y \tau_3'(\eta) N(1, y; 0, \eta) d\eta + \\ &+ \int_0^y \varphi_1'(\eta) N(1, y; 1, \eta) d\eta + \int_0^1 \tau_1'(\xi) N(1, y; \xi, 0) d\xi + \\ &+ \int_0^y e^{c\eta} d\eta \int_0^{1-\eta} [\tau_1''(\xi + \eta) - \nu_1(\xi + \eta)] N_\xi(1, y; \xi, \eta) d\xi + \\ &+ \int_0^y e^{c\eta} d\eta \int_{1-\eta}^1 \omega_{12}(\xi + \eta) N_\xi(1, y; \xi, \eta) d\xi. \end{aligned} \tag{35}$$

Here and at the top of the functions $G(x, y; \xi, \eta)$ and $N(x, y; \xi, \eta)$ have the form:

$$\left. \begin{matrix} G(x, y; \xi, \eta) \\ N(x, y; \xi, \eta) \end{matrix} \right\} = \frac{1}{2\sqrt{\pi(y-\eta)}} \sum_{n=-\infty}^{+\infty} \left\{ \exp \left[-\frac{(x-\xi-2n)^2}{4(y-\eta)} \right] \mp \exp \left[-\frac{(x+\xi-2n)^2}{4(y-\eta)} \right] \right\}.$$

They are Green's functions of the first and second boundary value problems for the heat equation. Substituting (32) into (33), after some transformations, we have the equation

$$\tau_3'(y) + \int_0^y \tau_3'(\eta)K_1(y, \eta)d\eta + \int_0^y K_2(y, \eta)\omega_{12}(1 + \eta)d\eta = g_1(y). \quad (36)$$

And differentiating the equation (35) after some calculations, we obtain the Volterra integral equation of the second kind with respect to $\omega_{12}(1 + y)$:

$$\omega_{12}(1 + y) + \int_0^y K_3(y, \eta)\omega_{12}(1 + \eta)d\eta + \int_0^y \tau_3'(\eta)K_4(y, \eta)d\eta = g_2(y), \quad (37)$$

where $K_1(y, \eta)$, $K_2(y, \eta)$, $K_3(y, \eta)$, $K_4(y, \eta)$, $g_1(y)$, $g_2(y)$ are known functions, and $K_1(y, \eta)$, $K_3(y, \eta)$ have a weak singularity ($\frac{1}{2}$), and $K_2(y, \eta)$, $K_4(y, \eta)$, $g_1(y)$, $g_2(y)$ are continuous functions.

Solving the system of equations (36), (37), we find the functions $\tau_3'(y)$, $\omega_{12}(1 + y)$ and thus, the functions $\nu_3(y)$, $\omega_{32}(y)$, $u_1(x, y)$, $u_3(x, y)$.

Remark 1. The case when $-1 < \gamma < 0$, the problem is investigated by dividing the domain G_1 into n parts whose heights of the first $n - 1$ domains are equal to $-\frac{b}{a}$, and the last – no more than $-\frac{b}{a}$. The problem is solved in each domain sequentially, similar to the case of $\gamma = -1$.

Remark 2. In [4, 11], a number of boundary value problems for more general equations of the third and fourth orders of parabolic-hyperbolic type in a domain with a single line of type change were considered.

Conclusion

This work presents the formulation and comprehensive analysis of a boundary value problem for a third-order parabolic-hyperbolic PDE within a geometrically intricate pentagonal domain. Through the development of a constructive method for the equation's solution, we have established its unique solvability. Our findings enrich the theoretical underpinnings of mixed-type equations and extend the toolkit for addressing boundary value problems in domains with complex geometries. This research not only advances our understanding of parabolic-hyperbolic equations of third order but also has potential implications for their application in modeling multifaceted physical systems and phenomena. Future studies may explore the application of these findings in practical scenarios and the investigation of similar problems in higher-dimensional spaces or with more complex boundary conditions.

Author Contributions

All authors contributed equally to this work.

Conflict of Interest

The authors declare no conflict of interest.

References

- 1 Гельфанд И.М. Некоторые вопросы анализа и дифференциальных уравнений / И.М. Гельфанд // Успехи математических наук. — 1959. — Т. 14. Вып. 3(87). — С. 3–19.
- 2 Джураев Т.Д. Краевые задачи для уравнений параболо-гиперболического типа / Т.Д. Джураев, А. Сопуев, М. Мамажанов. — Ташкент: Фан, 1986. — 220 с.

- 3 Джураев Т.Д. Краевые задачи для одного класса уравнений четвертого порядка смешанного типа / Т.Д. Джураев, М. Мамажанов // Дифференциальные уравнения. — 1986. — Т. 22, № 1. — С. 25–31.
- 4 Мамажанов М. Постановка и метод решения некоторых краевых задач для одного класса уравнений третьего порядка парабола-гиперболического типа / М. Мамажанов, Х. Шерматова, Х. Мукаддасов // Вестн. КРАУНЦ. Физ.-мат. науки. — 2014. — № 1(8). — С. 7–13.
- 5 Мамажонов М. Постановка и метод исследования некоторых краевых задач для одного класса уравнений четвертого порядка парабола-гиперболического типа / М. Мамажонов, С.М. Мамажонов // Вестн. КРАУНЦ. Физ.-мат. науки. — 2014. — № 1(8). — С. 14–19.
- 6 Апаков Ю.П. Краевая задача для неоднородного уравнения четвертого порядка с младшими членами / Ю.П. Апаков, С.М. Мамажанов // Дифференциальные уравнения. — 2023. — Т. 59, № 2. — С. 183–192. <https://doi.org/10.31857/S037406412302005X>
- 7 Apakov Y.P. Boundary Value Problem for a Fourth-Order Equation of the Parabolic-Hyperbolic Type with Multiple Characteristics with Slopes Greater Than One / Y.P. Apakov, S.M. Mamajonov // Russ Math. — 2022 — Vol. 66. — P. 1–11. <https://doi.org/10.3103/S1066369X22040016>
- 8 Apakov Yu.P. Solvability of a Boundary Value Problem for a Fourth Order Equation of Parabolic-Hyperbolic Type in a Pentagonal Domain / Yu.P. Apakov, S.M. Mamajonov // Journal of Applied and Industrial Mathematics. — 2021. — Vol. 15. — P. 586–596. <https://doi.org/10.1134/S1990478921040025>
- 9 Мамажонов М. Об одной краевой задаче для уравнения третьего порядка парабола-гиперболического типа в вогнутой шестиугольной области / М. Мамажонов, Х.М. Шерматова // Вестн. КРАУНЦ. Физ.-мат. науки. — 2017. — № 1(17). — С. 14–21. <https://doi.org/10.18454/2079-6641-2017-17-1-14-21>
- 10 Shermatova K.M. Investigation of a boundary-value problem for a third order parabolic hyperbolic equation in the form $(b\frac{\partial}{\partial y} + c)(Lu) = 0$ / K.M. Shermatova // Theoretical Applied Science. — 2020. — Vol. 87, Iss. 07. — P. 160–165. <https://dx.doi.org/10.15863/TAS.2020.07.87.37>
- 11 Мамажонов М. Об одной краевой задаче для уравнения парабола-гиперболического типа третьего порядка в вогнутой шестиугольной области / М. Мамажонов, Х.М. Шерматова, Т.Н. Мухторова // XIII Белорус. мат. конф.: матер. Междунар. науч. конф. (22–25 ноября 2021 года). — Минск, 2021. — С. 67–68.
- 12 Mamajonov M. On a boundary value problem for a third-order parabolic-hyperbolic equation in a pentagonal domain with three lines of type change, whose hyperbolic parts are triangles / M. Mamajonov, K. Shermatova, O. Makhkamova // International Journal of social science & Interdisciplinary Research. — 2022. — Vol. 11. — P. 111–116.
- 13 Mamajonov M. On a Boundary Value Problem for a Third-Order Equation of the Parabolic-Hyperbolic Type in a Triangular Domain with Three Type Change Lines / M. Mamajonov, H.M. Shermatova // Journal of Applied and Industrial Mathematics. — 2022. — Vol. 16. — P. 481–489. <https://doi.org/10.1134/S1990478922030127>
- 14 Mamajonov M. Statement and study of a boundary value problem for a third-order equation of parabolic-hyperbolic type in a mixed pentagonal domain, when the slope of the characteristic of the operator the first order is greater than one / M. Mamajonov, X. Shermatova // International journal of research in commerce, IT, engineering and social sciences. — 2022. — Vol. 16, No. 5. — P. 117–130.
- 15 Shermatova Kh.M. Investigation of a boundary value problem for a third order parabolic hyperbolic equation in the form $(b\frac{\partial}{\partial y} + c)(Lu) = 0$ / Kh.M. Shermatova // Scientific Bulletin of Namangan State University. — 2020. — No. 2(4). — P. 44–53.

Параболалық-гиперболалық типтегі үшінші ретті теңдеу үшін шекаралық шама есебін тұжырымдау және зерттеу туралы

М. Мамажонов¹, К. Рахимов², Х. Шерматова²

¹Қоқан мемлекеттік педагогикалық институты, Қоқан, Өзбекстан;

²Фергана мемлекеттік университеті, Фергана, Өзбекстан

Мақалада параболалық және гиперболалық аймақтардан тұратын бесбұрышты аймақтағы параболалық-гиперболалық типтегі үшінші ретті ішінара туындылардағы теңдеудің жаңа шекті есебі зерттелген. Мұндай теңдеулер физика, инженерия және қаржы сияқты әртүрлі салалардағы күрделі физикалық құбылыстарды модельдеуде шешуші рөл атқарады, олардың аралас типтегі табиғатына орай динамиканың кең ауқымын инкапсуляциялау қабілетіне байланысты. Шешудің конструктивті тәсілін қолдана отырып, біз қойылған есептің біржақты шешілуін көрсетеміз. Бұл зерттеудің маңыздылығы математикалық физика мен сабақтас пәндердегі теориялық және қолданбалы зерттеулерге жаңа мүмкіндіктер ашатын күрделі геометриялық салалардағы жоғары ретті жартылай дифференциалдық теңдеулерді түсіну және шешу үшін математикалық негізді кеңейту болып табылады.

Клт сөздер: параболалық-гиперболалық типтегі дифференциалдық теңдеулер, үшінші ретті параболалық-гиперболалық тип.

О постановке и исследовании краевой задачи для уравнения третьего порядка параболо-гиперболического типа

М. Мамажанов¹, К. Рахимов², Х. Шерматова²

¹Кокандский государственный педагогический институт, Коканд, Узбекистан;

²Ферганский государственный университет, Фергана, Узбекистан

В статье исследована новая краевая задача для уравнения в частных производных третьего порядка параболо-гиперболического типа в пятиугольной области, состоящей как из параболических, так и из гиперболических областей. Такие уравнения играют решающую роль в моделировании сложных физических явлений в различных областях, таких как физика, инженерия и финансы, благодаря их способности инкапсулировать широкий диапазон динамики из-за своей природы смешанного типа. Используя конструктивный подход к решению, мы демонстрируем однозначную разрешимость поставленной задачи. Значимость этого исследования заключается в расширении математической основы для понимания и решения смешанных уравнений в частных производных высшего порядка в сложных геометрических областях, что открывает новые возможности для теоретических и прикладных исследований в математической физике и смежных дисциплинах.

Ключевые слова: дифференциальные уравнения параболо-гиперболического типа, параболо-гиперболический тип третьего порядка.

References

- 1 Gelfand, I.M. (1959). Nekotorye voprosy analiza i differentsialnykh uravnenii [Some issues of analysis and differential equations]. *Uspekhi matematicheskikh nauk*, 14, 3(87), 3–19 [in Russian].
- 2 Dzhuraev, T.D., Sopuev, A., & Mamazhanov, M. (1986). *Kraevye zadachi dlia uravnenii parabologiperbolicheskogo tipa* [Boundary value problems for parabolic-hyperbolic equations]. Tashkent: Fan [in Russian].
- 3 Dzhuraev, T.D. & Mamazhanov, M. (1986). Kraevye zadachi dlia odnogo klassa uravnenii chetvertogo poriadka smeshannogo tipa [Boundary value problems for a class of fourth-order mixed-type equations]. *Differentsialnye uravneniia – Differential Equations*, 22(1), 25–31 [in Russian].

- 4 Mamazhanov, M., Shermatova, Kh., & Mukaddasov, Kh. (2014). Postanovka i metod resheniia nekotorykh kraevykh zadach dlia odnogo klassa uravnenii tretego poriadka parabolo-giperbolicheskogo tipa [Formulation and method of solving some boundary value problems for a class of third-order equations of parabolic-hyperbolic type]. *Vestnik KRAUNTs. Fiziko-matematicheskie nauki — Bulletin of Kraesc. Physical and Mathematical Sciences*, 1(8), 7–13 [in Russian].
- 5 Mamajonov, M., & Mamajonov, S.M. (2014). Postanovka i metod issledovaniia nekotorykh kraevykh zadach dlia odnogo klassa uravnenii chetvertogo poriadka parabolo-giperbolicheskogo tipa [Formulation and method of investigation of some boundary value problems for a class of fourth-order equations of parabolic-hyperbolic type]. *Vestnik KRAUNTs. Fiziko-matematicheskie nauki — Bulletin of Kraesc. Physical and Mathematical Sciences*, 1(8), 14–19 [in Russian].
- 6 Apakov, Yu.P., & Mamazhonov, S.M. (2023). Kraevaia zadacha dlia neodnorodnogo uravneniia chetvertogo poriadka s mladshimi chlenami [A boundary value problem for an inhomogeneous fourth-order equation with minor terms]. *Differentsialnye uravneniia — Differential equations*, 59(2), 183–192 [in Russian]. <https://doi.org/10.31857/S037406412302005X>
- 7 Apakov, Y.P., & Mamajonov, S.M. (2022). Boundary Value Problem for a Fourth-Order Equation of the Parabolic-Hyperbolic Type with Multiple Characteristics with Slopes Greater Than One. *Russ Math.*, 66, 1–11. <https://doi.org/10.3103/S1066369X22040016>
- 8 Apakov, Y.P., & Mamajonov, S.M. (2021). Solvability of one boundary value problem for a fourth-order equation of parabolic-hyperbolic type in a pentagonal domain. *Journal of Applied and Industrial Mathematics*, 24(4), 25–38. <https://doi.org/10.1134/S1990478921040025>
- 9 Mamazhonov, M., & Shermatova, Kh.M. (2017). Ob odnoi kraevoi zadache dlia uravneniia tretego poriadka parabolo-giperbolicheskogo tipa v vognutoi shestiugolnoi oblasti [On a boundary value problem for a third-order equation of parabolic-hyperbolic type in a concave hexagonal domain] *Vestnik KRAUNTs. Fiziko-matematicheskie nauki — Bulletin of Kraesc. Physical and Mathematical Sciences*, 1(17), 14–21 [in Russian]. <https://doi.org/10.18454/2079-6641-2017-17-1-14-21>
- 10 Shermatova, K.M. (2020). Investigation of a boundary-value problem for a third order parabolic hyperbolic equation in the form $(b\frac{\partial}{\partial y} + c)(Lu) = 0$. *Theoretical Applied Science*, 87(7), 160–165.
- 11 Mamajonov, M., Shermatova, Kh.M., & Mukhtorova, T.N. (2021). Ob odnoi kraevoi zadache dlia uravneniia parabolo-giperbolicheskogo tipa tretego poriadka v vognutoi shestiugolnoi oblasti [On a boundary value problem for a third-order parabolic-hyperbolic equation in a concave hexagonal domain]. *XIII Belorusskaia matematicheskaiia konferentsiia: materialy Mezhdunarodnoi nauchnoi konferentsii — XIII Belarusian Mathematical Conference: Proceedings of the International Scientific Conference (22–25 noiabria 2021 goda)*. Minsk, 67–68 [in Russian].
- 12 Mamajonov, M., Shermatova, Kh.M., & Makhkamova, O. (2022). On a boundary value problem for a third-order parabolic-hyperbolic equation in a pentagonal domain with three lines of type change, whose hyperbolic parts are triangles. *International Journal of social science & Interdisciplinary Research*, 11, 111–116.
- 13 Mamajonov, M., & Shermatova, Kh.M. (2022). On a Boundary Value Problem for a Third-Order Equation of the Parabolic–Hyperbolic Type in a Triangular Domain with Three Type Change Lines. *Journal of Applied and Industrial Mathematics*, 16, 481–489.
- 14 Mamajonov, M., Shermatova, Kh. (2022). Statement and study of a boundary value problem for a third-order equation of parabolic-hyperbolic type in a mixed pentagonal domain, when the slope of the characteristic of the operator the first order is greater than one. *International journal of research in commerce, IT, engineering and social sciences*, 16(5), 117–130.
- 15 Shermatova, Kh.M. (2020). Investigation of a boundary value problem for a third order parabolic hyperbolic equation in the form $(b\frac{\partial}{\partial y} + c)(Lu) = 0$. *Scientific Bulletin of Namangan State University*, 2(4), 44–53.

*Author Information**

Mirza Mamajonov — Candidate of physical and mathematical sciences, docent, Associate Professor of the Department of Mathematics, Kokand State Pedagogical Institute, 23, Turon street, Kokand, Uzbekistan; e-mail: mirzamamajonov@gmail.com; <https://orcid.org/0009-0003-8413-0549>

Quvvatali Ortikovich Rakhimov (*corresponding author*) — Doctor of philosophy in technical sciences, docent, Head of the Department of Information Technology, Fergana State University, 19, Murabbiylar street, Fergana, Uzbekistan; e-mail: quvvatali.rahimov@gmail.com; <https://orcid.org/0000-0002-1863-3645>

Khilolaxon Mirzayevna Shermatova — Senior Lecturer at the Department of Information Technology, Fergana State University, 19, Murabbiylar street, Fergana, Uzbekistan; e-mail: shermatovahilola1978@gmail.com; <https://orcid.org/0000-0001-5014-9549>

*The author's name is presented in the order: First, Middle and Last Names.